ELSEVIER

Environmental Pollution 142 (2006) I–V

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 142

Number 1 2006

1 **Urban environmental geochemistry of trace metals**
C.S.C. Wong, X. Li, I. Thornton

Urban environmental geochemistry as a scientific discipline provides valuable information on trace metal contamination of the urban environment and its associated health effects.

17 **Assessment of heavy metal bioavailability in contaminated sediments and soils using green fluorescent protein-based bacterial biosensors**
V.H.-C. Liao, M.-T. Chien, Y.-Y. Tseng, K.-L. Ou

Nonpathogenic GFP-based bacterial biosensor is applicable in determining the bioavailability of heavy metals in environmental samples.

24 **Effects of oil sands tailings compounds and harsh weather on mortality rates, growth and detoxification efforts in nestling tree swallows (*Tachycineta bicolor*)**
M.-L. Gentes, C. Waldner, Z. Papp, J.E.G. Smits

Under natural stress caused by harsh weather, birds exposed to chemicals from the oil sands extraction process suffered higher mortality than those in control areas.

34 **Ectomycorrhizas impede phytoremediation of polycyclic aromatic hydrocarbons (PAHs) both within and beyond the rhizosphere**
E.J. Joner, C. Leyval, J.V. Colpaert

The ectomycorrhizal fungus S. bovinus impeded degradation of PAHs in soil, probably due to its negative effect on the availability of mineral nutrients of more potent PAH degraders.

39 **Characterization of trace organic contaminants in marine sediment from Yeongil Bay, Korea: 1. Instrumental analyses**
C.-H. Koh, J.S. Khim, D.L. Villeneuve, K. Kannan, J.P. Giesy

Among various sediment contaminant classes measured, nonylphenol and PAHs are responsible for the variability among sampling sites, suggesting the existence of multiple sources in Yeongil Bay sediment.

48 **Characterization of trace organic contaminants in marine sediment from Yeongil Bay, Korea: 2. Dioxin-like and estrogenic activities**
C.-H. Koh, J.S. Khim, D.L. Villeneuve, K. Kannan, J.P. Giesy

In vitro bioassay responses observed for Yeongil Bay surficial sediment and sediment core extracts showed the greatest dioxin-like and estrogenic activities in the mid-polarity fraction containing PAHs as well as chlorinated dioxins and furans.

58 **Effects of NO_2 and NH_3 from road traffic on epiphytic lichens**
L. Frati, E. Caprasecca, S. Santoni, C. Gaggi, A. Guttova, S. Gaudino, A. Pati, S. Rosamilia, S.A. Pirintsos, S. Loppi

NH_3 and NO_2 from a busy road in central Italy do not influence epiphytic lichens.

65 **Genotoxicity biomarkers and acetylcholinesterase activity in natural populations of *Mytilus galloprovincialis* along a pollution gradient in the Gulf of Oristano (Sardinia, western Mediterranean)**
P. Magni, G. De Falco, C. Falugi, M. Franzoni, M. Monteverde, E. Perrone, M. Sgro, C. Bolognesi

Biomarkers can be useful in coastal marine biomonitoring.

73 ***Brachidontes variabilis* and *Patella* sp. as quantitative biological indicators for cadmium, lead and mercury in the Lebanese coastal waters**
K.F. Nakhlé, D. Cossa, G. Khalaf, B. Beliaeff

Two molluscs were effective bioindicators for metal pollution in waters along the Lebanese coast.

83 **California sea lions (*Zalophus californianus californianus*) have lower chlorinated hydrocarbon contents in northern Baja California, México, than in California, USA**
L.D. Toro, G. Heckel, V.F. Camacho-Ibar, Y. Schramm

Results suggest that sea lion prey must also have lower hydrocarbons in Baja California than in California in the USA.

93 Lead shot from hunting as a source of lead in human blood
P. Johansen, H.S. Pedersen, G. Asmund, F. Riget

Birds hunted with lead shot and consumed are a source of lead in human blood.

98 Imposex occurrence in marine whelks at a military facility in the high Arctic
J. Strand, C.M. Glahder, G. Asmund

Imposex as a biomarker was more sensitive for TBT than analytical chemical methods.

103 Bioavailability and cellular effects of metals on *Lumbricus terrestris* inhabiting volcanic soils
A. Amaral, M. Soto, R. Cunha, I. Marigómez, A. Rodrigues

In earthworms, differences in the chloragogenous tissue morphometry may be related to the bioavailability of metals in soils.

109 Growth and nutritive quality of *Poa pratensis* as influenced by ozone and competition
J. Bender, R.B. Muntifering, J.C. Lin, H.J. Weigel

Early-season O_3 exposure decreased nutritive quality of Poa pratensis, and nutritive quality response to O_3 was not altered by interspecific competition.

116 Effects of metals on condition and reproductive output of the smooth toadfish in Sydney estuaries, south-eastern Australia
R. Alquezar, S.J. Markich, D.J. Booth

Enhanced metal levels in sediment reduced reproductive output and condition in toadfish.

123 Metal accumulation in the smooth toadfish, *Tetractenos glaber*, in estuaries around Sydney, Australia
R. Alquezar, S.J. Markich, D.J. Booth

Metal levels in toadfish tissues reflect sediment metal levels and show gender differences.

132 Sorption and bioreduction of hexavalent uranium at a military facility by the Chesapeake Bay
W. Dong, G. Xie, T.R. Miller, M.P. Franklin, T.P. Oxenberg, E.J. Bouwer, W.P. Ball, R.U. Halden

At the Aberdeen Proving Ground in Maryland, USA, migration of depleted uranium into the Chesapeake Bay is limited by rapid sorption of the radionuclide to natural organic matter followed by slow biological reduction of water-soluble U(VI) to the insoluble and less toxic U(IV) species.

143 Air-vegetation transfer of PCDD/PCDFs: An assessment of field data and implications for modeling
M. Schuhmacher, K.C. Jones, J.L. Domingo

Soil and air transfer of PCDD/PCDFs to vegetation were evaluated in residential and industrial areas.

151 Comparison of the role of the sea club-rush *Scirpus maritimus* and the sea rush *Juncus maritimus* in terms of concentration, speciation and bioaccumulation of metals in the estuarine sediment
C.M.R. Almeida, A.P. Mucha, M.T.S.D. Vasconcelos

Both Juncus maritimus and Scirpus maritimus have potential for the phytostabilization of Cd, and the last one also for Pb phytostabilization.

160 Fate of heavy metals after application of sewage sludge and wood–ash mixtures to short-rotation willow coppice
I. Dimitriou, J. Eriksson, A. Adler, P. Aronsson, T. Verwijst

Commercially applied, legally regulated applications of sewage sludge and wood–ash mixtures to short-rotation willow coppice in the Enköping area, central Sweden, are tested.

170 Atmospheric concentrations and air–sea exchanges of nonylphenol, tertiary octylphenol and nonylphenol monoethoxylate in the North Sea
Z. Xie, S. Lakaschus, R. Ebinghaus, A. Caba, W. Ruck

Air–sea vapour exchanges of estrogenic alkylphenols are studied in the North Sea and it is shown that a net deposition is occurring.

181 Aquatic risk assessment of the new rice herbicide profoxydim
P. Sánchez, J. Kubitza, G. Peter Dohmen, J.V. Tarazona

Risk refinement assessment of rice pesticides starting with DTA and moving to community studies is a cost-effective approach, only if required.

Number 2 2006

191 Biodegradation and speciation of residual SS-ethylenediaminedisuccinic acid (EDDS) in soil solution left after soil washing
S. Tandy, A. Ammann, R. Schulin, B. Nowack

Even in polluted soils, EDDS is degraded.

200 Use of hydra for chronic toxicity assessment of waters intended for human consumption
V.V. Arkhipchuk, C. Blaise, M.V. Malinovskaya

Hydra is a reliable and relevant test organism for the assessment of acute and chronic toxicity of waters intended for human consumption.

212 Consequences for *Protaphorura armata* (Collembola: Onychiuridae) following exposure to genetically modified *Bacillus thuringiensis* (*Bt*) maize and non-*Bt* maize
L.-H. Heckmann, B.S. Griffiths, S. Caul, J. Thompson, M. Pusztai-Carey, W.J. Moar, M.N. Andersen, P.H. Krogh

Protaphorura armata performed equally well when reared on two Bt and three non-Bt maize varieties.

217 Physicochemical characterization and mercury speciation of particle-size soil fractions from an abandoned mining area in Mieres, Asturias (Spain)
R. Fernández-Martínez, J. Loredo, A. Ordóñez, M.I. Rucandio

Factors affecting mercury mobility in soils influenced by abandoned mining.

227 A spatially-evaluated methodology for assessing risk to a population from contaminated land
J.R. Gay, A. Korre

A methodology for calculating and mapping risks to a population from intake of soil contaminants.

235 Polychlorinated biphenyl congener patterns in tree swallows (*Tachycineta bicolor*) nesting in the Housatonic River watershed, western Massachusetts, USA, using a novel statistical approach
C.M. Custer, L.B. Read

Polychlorinated biphenyl congener patterns in swallows differed between the main stem of the Housatonic River, MA and other locations in the watershed.

246 Impact of short-term and long-term elevated CO_2 on emission of carbonyls from adult *Quercus petraea* and *Carpinus betulus* trees
J. Kreuzwieser, H. Rennenberg, R. Steinbrecher

Carbonyl exchange of European tree species was not affected by elevated ambient carbon dioxide concentrations.

254 Spatial variability and temporal stability of throughfall deposition under beech (*Fagus sylvatica* L.) in relationship to canopy structure
J. Staelens, A. De Schrijver, K. Verheyen, N.E.C. Verhoest

Ion deposition under a deciduous beech tree was strongly affected by the canopy structure throughout the entire year.

264 The marine macroalga *Cystoseira baccata* as biosorbent for cadmium(II) and lead(II) removal: Kinetic and equilibrium studies
P. Lodeiro, J.L. Barriada, R. Herrero, M.E. Sastre de Vicente

Marine macroalgae show promise for biosorption of lead and cadmium.

274 Benthic foraminifera and heavy metals distribution: A case study from the Naples Harbour (Tyrrhenian Sea, Southern Italy)
L. Ferraro, M. Sprovieri, I. Alberico, F. Lirer, L. Prevedello, E. Marsella

Benthic foraminiferal density and species distribution may be used as pollution indicators.

288 Influence of vegetation in mitigation of methyl parathion runoff
M.T. Moore, E.R. Bennett, C.M. Cooper, S. Smith Jr., J.L. Farris, K.G. Drouillard, R. Schulz

Vegetated wetlands are three times more effective at mitigating methyl parathion runoff than are non-vegetated wetlands.

295 Evaluating the vulnerability of surface waters to antibiotic contamination from varying wastewater treatment plant discharges
A.L. Batt, I.B. Bruce, D.S. Aga

This work investigates the extent of antibiotic concentrations in receiving waters from discharges of wastewater treatment plants.

303 Metal stoichiometry in predicting Cd and Cu toxicity to a freshwater green alga *Chlamydomonas reinhardtii*
W.-X. Wang, R.C.H. Dei

The stoichiometry of metals and P can be used to predict metal toxicity to freshwater phytoplankton under different P conditions.

313 Seasonal variations in hepatic Cd and Cu concentrations and in the sub-cellular distribution of these metals in juvenile yellow perch (*Perca flavescens*)
L.D. Kraemer, P.G.C. Campbell, L. Hare

In fish from metal-contaminated sites, seasonal variations in hepatic Cd and Cu concentrations were greater than in fish from reference sites, and homeostatic control of sub-cellular metal distribution was compromised.

326 Electrokinetic enhancement of phenanthrene biodegradation in creosote-polluted clay soil
J.-L. Niqui-Arroyo, M. Bueno-Montes, R. Posada-Baquero, J.-J. Ortega-Calvo

Electrokinetic bioremediation is a potentially effective technology to treat PAH-polluted, clay-rich soils.

333 Calibration of the Chemcatcher passive sampler for the monitoring of priority organic pollutants in water
B. Vrana, G.A. Mills, E. Dominiak, R. Greenwood

A calibration method and data that are required for in situ measurement of the time-weighted average concentrations of hydrophobic priority organic pollutants in water using Chemcatcher passive sampling device are presented.

344 Herbicide contamination of surficial groundwater in Northern Italy
L. Guzzella, F. Pozzoni, G. Giuliano

Herbicide groundwater pollution is an environmental problem in Northern Italy.

354 Field responses of *Prunus serotina* and *Asclepias syriaca* to ozone around southern Lake Michigan
J.P. Bennett, E.A. Jepsen, J.A. Roth

Decreased cherry branch elongation, milkweed stem height and pod production, and foliar injury on both species occurred at sites around southern Lake Michigan at ozone exposures of 13 SUM06 ppm-h and 93–98 ppb peak hourly.

367 Perfluorooctanesulfonate and related fluorochemicals in biological samples from the north coast of Colombia
J. Olivero-Verbel, L. Tao, B. Johnson-Restrepo, J. Guette-Fernández, R. Baldiris-Avila, I. O'byrne-Hoyos, K. Kannan

Perfluorooctanesulfonate and related perfluorinated compounds have been found in a tropical ecosystem of South America.

373 Heavy metals affect the coelomocyte-bacteria balance in earthworms: Environmental interactions between abiotic and biotic stressors
E. Olchawa, M. Bzowska, S.R. Stürzenbaum, A.J. Morgan, B. Plytycz

Metals affect coelomocyte-bacterial balance in earthworms.

Number 3 2006

383 Reproductive transfer of organochlorines in viviparous surfperch, *Ditrema temmincki*
M. Oka, T. Arai, Y. Shibata, N. Miyazaki

The reproductive transfer of POPs in matrotrophic viviparous surfperch was determined for the first time.

388 Atmospheric polycyclic aromatic hydrocarbons (PAHs) in Asia: A review from 1999 to 2004
K.-F. Chang, G.-C. Fang, J.-C. Chen, Y.-S. Wu

This work summarizes the characteristics, sources and distribution, and the healthy impacts of atmospheric PAH species in Asia.

397 Selenium and other trace metals in fish inhabiting a fly ash stream: Implications for regulatory tissue thresholds
R.J. Reash, T.W. Lohner, K.V. Wood

Bluegill sunfish inhabiting a coal fly ash receiving stream had elevated selenium levels in whole body and gonad tissue (9–10 times higher than reference fish), and antagonistic metal interactions may be one of several mechanisms allowing long-term persistence of the population.

409 Geochemical and Pb isotopic evidence for sources and dispersal of metal contamination in stream sediments from the mining and smelting district of Příbram, Czech Republic
V. Ettler, M. Mihaljevič, O. Šebek, M. Molek, T. Grygar, J. Zeman

Pb isotopes properly complete traditional investigations of metal sources and dispersal in contaminated stream sediments.

418 Mercury accumulation in surface sediments of salt marshes of the Bay of Fundy
G.A. Hung, G.L. Chmura

Short-term storage of mercury was measured in salt marsh sediments of the Bay of Fundy.

432 Eco-toxicological studies of diesel and biodiesel fuels in aerated soil
A. Lapinskienė, P. Martinkus, V. Rėbždaitė

The comparison of diesel and biodiesel fuels' eco-toxicological parameters in non-adapted aerated soil is relevant when considering the effects of these substances on the environment in cases of accidental spills.

438 Non-destructive pollution exposure assessment in the European hedgehog (*Erinaceus europaeus*): II. Hair and spines as indicators of endogenous metal and As concentrations
H. D'Havé, J. Scheirs, V.K. Mubiana, R. Verhagen, R. Blust, W. De Coen

Hedgehog hair and spines are promising non-destructive biomonitoring tools of terrestrial metal pollution.

449 Cosorption study of organic pollutants and dissolved organic matter in a soil
F. Flores-Céspedes, M. Fernández-Pérez, M. Villafranca-Sánchez, E. González-Pradas

Cosorption of organic pollutants and DOM.

457 High incidence of deformity in aquatic turtles in the John Heinz National Wildlife Refuge
B. Bell, J.R. Spotila, J. Congdon

This paper presents findings on the prevalence of developmental abnormalities in turtles at a national wildlife refuge that have direct relevance to studies on the effects of contamination on development and morphology of vertebrates.

466 Polycyclic aromatic hydrocarbons affect survival and development of common snapping turtle (*Chelydra serpentina*) embryos and hatchlings
R.J. Van Meter, J.R. Spotila, H.W. Avery

Exposure to polycyclic aromatic hydrocarbons on the egg reduces survival of snapping turtle embryos and causes developmental abnormalities.

476 Tissue-specific accumulation and lactational transfer of polychlorinated biphenyls, chlorinated pesticides, and brominated flame retardants in hooded seals (*Cistophora cristata*) from the Gulf of St. Lawrence: Applications for monitoring
H. Wolkers, M.O. Hammill, B. van Bavel

Blood plasma contaminant concentrations differ from those in other tissues and might therefore be unsuitable for contaminant monitoring in hooded seals.

487 Presence of plutonium contamination in soils from Palomares (Spain)
M.C. Jiménez-Ramos, R. García-Tenorio, I. Vioque, G. Manjón, M. García-León

The remaining transuranic contamination in Palomares soils is mainly present in particulate form.

493 Linking biosensor responses to Cd, Cu and Zn partitioning in soils
J.J.C. Dawson, C.D. Campbell, W. Towers, C.M. Cameron, G.I. Paton

Biosensors link biological hazard assessments of metals in soils with physico-chemical partitioning.

501 Using multivariate analyses and GIS to identify pollutants and their spatial patterns in urban soils in Galway, Ireland
C. Zhang

Multivariate statistics and GIS are useful tools to identify pollutants in urban soils.

512 Characterization of contamination in and toxicities of a shipyard area in Hong Kong
S.W. Chiu, K.M. Ho, S.S. Chan, O.M. So, K.H. Lai

Heavily contaminated shipyards result in high ecotoxicities but the environmental conditions handicap natural degradation.

521 Study of the trace metal ion influence on the turnover of soil organic matter in cultivated contaminated soils
C. Dumat, K. Quenea, A. Bermond, S. Toinen, M.F. Benedetti

Measure of $\delta^{13}C$ and total metal concentrations in size fractions of contaminated soils suggests an influence of metals on the soil organic matter dynamic.

530 Field evaluation of in situ remediation of a heavy metal contaminated soil using lime and red-mud
C.W. Gray, S.J. Dunham, P.G. Dennis, F.J. Zhao, S.P. McGrath

Red mud was effective in immobilising heavy metals in soil.

540 Physiological and biochemical stress responses in grassland species are influenced by both early-season ozone exposure and interspecific competition
F. Scebba, F. Canaccini, A. Castagna, J. Bender, H.-J. Weigel, A. Ranieri

The competition between species modulates the ozone effect in a short-term.

549 Environmental contamination of mercury from Hg-mining areas in Wuchuan, northeastern Guizhou, China
G. Qiu, X. Feng, S. Wang, L. Shang

Mercury mining activities in Wuchun, Guizhou, China have resulted in seriously mercury contamination to the local environment.

559 Effects of eutrophication on vitellogenin gene expression in male fathead minnows (*Pimephales promelas*) exposed to 17α-ethynylestradiol in field mesocosms
D.A. Gordon, G.P. Toth, D.W. Graham, J.M. Lazorchak, T.V. Reddy, C.W. Knapp, F. deNoyelles Jr., S. Campbell, D.L. Lattier

Nutrient condition in surface waters strongly influences vitellogenin gene expression in male fathead minnows.

567 Geostatistical analysis and kriging of Hexachlorocyclohexane residues in topsoil from Tianjin, China
B.G. Li, J. Cao, W.X. Liu, W.R. Shen, X.J. Wang, S. Tao

Geostatistical analysis and kriging were applied to HCH concentrations in topsoil of Tianjin, China for mapping purposes.